

## **VYVGART HYTRULO®**

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

PATIENT INFORMATION (Complete or Fax Existing Chart)		PRESCRIBER INFORMATION	
Address:		Prescriber Name:	
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)			
Plan #:		Secondary Insurance (If Applicable):	
CLINICAL INFORMATION			
☐ G70.00 Myasthenia gravis without (acute) exacerbation (gMG) ☐ G70.01 Myasthenia gravis with (acute) exacerbation (gMG) ☐ G61.81 Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) ☐ Other: Anti-AChR Antibody Positive: ☐ Yes ☐ No MG-ADL score (if known): MGFA Classification (if known):			
ORDERS CONTROL OF THE			
Prescription type:  New start Restart Continued therapy Total Doses Received: Date of Last Injection/Infusion:			/Infusion:
Medication	•	Dose/Frequency Qty/Refills	
□ Vyvgart Hytrulo®	Myasthenia Gravis: Administer 1,008mg subcutaneously over 30 to 90 seconds once weekly Refills (treatment		Dispense Qty: Refills (treatment cycles):
Pre-Medication	Dose/Strength	Directions	
☐ Acetaminophen	□ 500mg	☐ Take 1-2 tablets PO prior to infusion or post-infusion as directed	
☐ Diphenhydramine	☐ 25mg IV/PO ☐ 50mg IV/PO	☐ Take 1 tablet PO prior to infusion or as directed OR ☐ Inject contents of 1 vial IV prior to infusion or as directed	
☐ Methylprednisolone	☐ 40mg ☐ 100mg ☐ 125mg	☐ Inject contents of 1 vial IV prior to infusion or as directed ☐ Other: Inject 100mg IV 30 minutes prior to infusion	
INFUSION REACTION ORDERS			
Mild reaction protocol:  ☑ Diphenhydramine 25mg IV, one time, for pruritus.  If symptoms worsen, see orders for moderate to severe reactions.  Moderate reaction protocol:  ☑ Acetaminophen 650mg PO, one time, for pyrexia or rigors			

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



## **VYVGART HYTRULO®**

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

Date: \_\_\_

Tunneled Catheter, and Noner: um 100 units/mL 3-5mL IV final als aline flush				
er: um 100 units/mL 3-5mL IV final Il saline flush				
er: um 100 units/mL 3-5mL IV final				
·				
$\boxtimes$ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw				
Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:				
Flushing Protocol (>66lbs/33kg)				
worsen				
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or				
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms				
□ Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis				
☐ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)				
If symptoms worsen, see interventions for severe reactions  Severe reaction protocol: (Call 911 if initiated):				
Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms				
☐ Diphenhydramine 50mg IV, one time, for pruritus or urticaria				
n aı				

**Prescriber Signature** 

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.