

## **BRIUMVI**<sup>TM</sup>

## Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)
Please Send a Copy of The Patient's Up-to-Date Clinical Notes

PATIENT INFORMATION (Complete or Fax Existing Chart)			PRESCRIBER INFORMATION						
Name:	ne: DOB:			Prescriber Name:					
Address:			State License:						
City, State, Zip:			NPI #: DEA:						
Phone: Alt. Phone:			Address:						
Email: SS#:			City, State, Zip:						
Gender:   M   F   Weight:(lbs)   Ht:			Phone: Fax:						
Allergies:			Office Contact: Phone:						
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)									
Primary Insurance:			Secondary Insurance (If Applicable):						
Plan #:			Plan #:						
Group #:			Group #:						
RX Card (PBM):			RX Card (PBM):						
BIN: PCN:	PCN:			BIN: F				PCN:	
CLINICAL INFORMATION									
☐ G35.A Relapsing-remitting ☐ G	G35.B0 Primary progressiv	☐ G35.B1 Active primary					☐ G35.B2 Non-active primary		
multiple sclerosis mul	tiple sclerosis, unspecified		progressive multiple sclerosis			clerosis	5	progressive multiple sclerosis	
☐ G35.C0 Secondary progressive ☐ G	G35.C1 Active secondary		$\square$ G35.CD Non-active secondary				$\square$ G35.D Multiple sclerosis,		
multiple sclerosis, unspecified pro	progressive multiple sclerosis			ssive multi	iple s	clerosis	unspecified		
☐ Other (Specify ICD-10 Code):									
Lab Orders: Frequency:									
Has patient received/plans on receiving any live or live-attenuated vaccinations 4 weeks prior to starting Briumvi™ treatment? ☐ Yes ☐ No									
Has Patient received/plans on receiving any non-live vaccinations 2 weeks prior to starting Briumvi™ treatment? ☐ Yes ☐ No									
Has Quantitative Serum Immunoglobulin Screening been performed?   Yes   No (Serum Immunoglobulin levels:)									
Has patient received an HBV Screening? ☐ Yes ☐ No (Results: ☐ Negative ☐ Positive)									
BRIUMVI™ ORDERS									
Prescription type: ☐ New start ☐ Restart ☐ Continued therapy Total I					Last Injection/Infusion:				
Medication		Dose/Frequency						Refills	
	☐ First Infusion: 150 mg (1 vial)								
☐ Briumvi™ 150mg vial	☐ Second Infusion: 450 mg (3 vials) (2 weeks after initial dose)						Refill:		
C	•	(3 vials) once every 24 weeks							
	Other:								
Pre-Medication	Route						Dose		
☐ Acetaminophen	□ PO	□ 500 —		☐ 650mg		1000n	ng		
☐ Methylprednisolone (Solu-Medrol)	□IV	☐ 60n	ng	□ 100 mg			mg		
☐ Diphenhydramine (Benadryl)	□ IV □ PO	☐ 25n	ng	□ 50mg					
Other:									
INFUSION REACTION ORDERS									
Mild reaction protocol:									
☐ Diphenhydramine 25mg IV, one time, for pruritus.									

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



will be verified upon insurance approval.

## **BRIUMVI**<sup>TM</sup>

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)
Please Send a Copy of The Patient's Up-to-Date Clinical Notes

Date: \_\_\_

Please Send a Copy of The Patient's Up-to-Date Clinical Notes If symptoms worsen, see orders for moderate to severe reactions. Moderate reaction protocol: ☐ Acetaminophen 650mg PO, one time, for pyrexia or rigors ☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria ☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms If symptoms worsen, see interventions for severe reactions Severe reaction protocol: (Call 911 if initiated): ☐ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%) ☑ Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis ☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen **FLUSHING & LOCKING ORDERS** Flushing Protocol (>66lbs/33kg) PIV and Midline: Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter: □ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion ☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw Locking Protocol (>66lbs/33kg) PICC: PIV and Midline: Implanted Port, Tunneled Catheter, and Nontunneled Catheter: □ Heparin Sodium 10 units/mL 1mL IV final ☐ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush flush post normal saline flush □ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush \*\* May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused **SIGNATURE** We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

Prescriber Signature

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.