



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Needs

PATIENT INFORMATION (Complete or Fax Existing Chart)		PRESCRIBER INFORMATION			
Name: DOB:		Prescriber Name:			
Address:		State License:	State License:		
City, State, Zip:		NPI #:Tax ID:			
Phone: Alt. Phone:		Address:			
Email: SS#:		City, State, Zip:			
Gender: M F Weight:(lbs) Ht:		Phone:			
Allergies:		Office Contact:	Phone:		
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)					
Primary Insurance:		Secondary Insurance (If Applicable):			
Plan #:		Plan #:			
Group #:		Group #:			
RX Card (PBM):		RX Card (PBM):			
BIN: PCN:		BIN: P			
CUNICAL INFORMATION					
CLINICAL INFORMATION					
☐ M32.0 Drug-induced Systemic Lupus Erythematosus ☐ M32.1 Systemic Lupus Erythematosus (organ or system involvement) ☐ L93.0 Discoid Lupus Erythematosus					
☐ M32.9 Systemic Lupus Erythematosus, unspecified ☐ Other:					
Has patient been previously treated for this condition? Yes ☐ No Is patient currently on therapy? ☐ Yes ☐ No					
BENLYSTA® ORDERS					
Prescription type: \square New start \square Restart \square Continued therapy Total Doses Received: Date of Last Injection/Infusion:					
Medication	Di	irections	Quantity/Refills		
	☐ 10mg/KG at 0, 2 and 4 weeks; then every 4 weeks		☐ Quantity:		
\square Benlysta $^{ ext{@}}$ (belimumab)	☐mg IV at 0, 2 and 4 weeks; then every 4 weeks ☐ Other:		☐ Refills:		
			□ neiii3		
INFUSION REACTION ORDERS					
Mild reaction protocol:					
oximes Diphenhydramine 25mg IV, one time, for	r pruritus.				
If symptoms worsen, see orders for moderate to severe reactions.					
Moderate reaction protocol:					
☑ Acetaminophen 650mg PO, one time, for pyrexia or rigors					
☐ Diphenhydramine 50mg IV, one time, for pruritus or urticaria					
☐ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms					
☑ Methylprednisolone 125mg IV, one time	r pruritus or urticaria	mptoms			
If symptoms worsen, see interventions for s	r pruritus or urticaria e, for respiratory or neurologic sy evere reactions	mptoms			
· · · · · · · · · · · · · · · · · · ·	r pruritus or urticaria e, for respiratory or neurologic sy evere reactions	mptoms			
If symptoms worsen, see interventions for s Severe reaction protocol: (Call 911 if initia ☑ Titrate oxygen via continuous flow per r	or pruritus or urticaria e, for respiratory or neurologic sy severe reactions ted): hasal cannula or face mask to mai	ntain spO2 of greater than ninety-five pe	rcent (>95%)		
If symptoms worsen, see interventions for some severe reaction protocol: (Call 911 if initia ☐ Titrate oxygen via continuous flow per row ☐ Diphenhydramine 50mg IV, one time, for	r pruritus or urticaria e, for respiratory or neurologic sy severe reactions ted): hasal cannula or face mask to mai r respiratory symptoms, edema, o	ntain spO2 of greater than ninety-five pe or anaphylaxis	rcent (>95%)		
If symptoms worsen, see interventions for some severe reaction protocol: (Call 911 if initia ☐ Titrate oxygen via continuous flow per roll ☐ Diphenhydramine 50mg IV, one time, for ☐ Methylprednisolone 125mg IV, one time	er pruritus or urticaria e, for respiratory or neurologic systewere reactions ted): lasal cannula or face mask to main respiratory symptoms, edema, ce, for respiratory symptoms, eder	ntain spO2 of greater than ninety-five pe or anaphylaxis na, or anaphylaxis	rcent (>95%)		
If symptoms worsen, see interventions for some severe reaction protocol: (Call 911 if initial ☐ Titrate oxygen via continuous flow per row Diphenhydramine 50mg IV, one time, for ☐ Methylprednisolone 125mg IV, one time ☐ Sodium Chloride 0.9% 500mL IV over 30	er pruritus or urticaria e, for respiratory or neurologic sy evere reactions ted): hasal cannula or face mask to mai r respiratory symptoms, edema, o e, for respiratory symptoms, eder -60 min, one time, for cardiovaso	ntain spO2 of greater than ninety-five pe or anaphylaxis ma, or anaphylaxis cular symptoms			
If symptoms worsen, see interventions for some severe reaction protocol: (Call 911 if initia ☐ Titrate oxygen via continuous flow per roll ☐ Diphenhydramine 50mg IV, one time, for ☐ Methylprednisolone 125mg IV, one time	er pruritus or urticaria e, for respiratory or neurologic sy evere reactions ted): hasal cannula or face mask to mai r respiratory symptoms, edema, o e, for respiratory symptoms, eder -60 min, one time, for cardiovaso	ntain spO2 of greater than ninety-five pe or anaphylaxis ma, or anaphylaxis cular symptoms			

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



ELLICHING & LOCKING OPDEDS



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Needs

FLOSTING & LOCKING ONDERS						
Flushing Protocol (>66lbs/33kg)						
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:				
\boxtimes 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion		☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mLIV flush after infusion/lab draw				
Locking Protocol (>66lbs/33kg)						
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-			
⋈ Heparin Sodium 10 units/mL 1mL IV final	☑ Heparin Sodium 10 units/mL 3mL IV final		tunneled Catheter:			
flush post normal saline flush	flush post normal saline flush		⋈ Heparin Sodium 100 units/mL 3-5mL IV final			
			flush post normal saline flush			
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused						
SIGNATURE						
We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.						
x			Date:			
Prescriber Signature						

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.