

REMICADE® (infliximab) & Biosimilar

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

DATIENT INFORMATION (C			DDECCRIDED INCOM	AATION	·	
PATIENT INFORMATION (Co			PRESCRIBER INFORM			
Name:	DOB:					
Address:			State License:			
City, State, Zip:					·	
Phone:	Alt. Phone:					
Email:	SS#:					
Gender: \square M \square F Weight: $_$	(lbs) Ht:				Fax:	
Allergies:			Office Contact:		Phone:	
INSURANCE INFORMATION	– AND – Send a copy of the	e patier	nt's prescription/insu	rance ca	rds (front & back)	
Primary Insurance:			Secondary Insurance (If A	pplicable)	:	
Plan #:						
Group #:						
RX Card (PBM):						
BIN:	PCN:				PCN:	
CLINICAL INFORMATION						
☐ K51.90 Moderate to Severe Ulc	erative Colitis		*If PPD test results are n	ot within 1	12 months inlease perform PPD	
☐ K50.90 Moderate to Severe Cro			*If PPD test results are not within 12 months, please perform PPD.			
☐ M06.9 Rheumatoid Arthritis	Jill 3 Disease		Tuberculosis Screening: PPD Test Date:			
☐ M45.9 Ankylosing Spondylitis			Results: Negative			
☐ L40.52 Psoriatic Arthritis			☐ Positive → ☐	Chest X-I	Ray Performed Date:	
			X-Ray Results: Negati	ive		
☐ L40.0 Plaque Psoriasis			\square Positive \Rightarrow TB treatment Initiated			
Other:						
Labs:		- CD			D 011	
		ESR q:		\ X-	Ray: □ Other:	
REMICADE®						
Prescription type: New start	☐ Restart ☐ Continued the	rapy	Total Doses Received:		Date of Last Infusion:	
Medication		Directio	ons		Quantity/Refills	
	Loading dose: 5mg/kg	m	a IV at week: 0.2.6		Loading dose: 3 doses. No refills.	
☐ Remicade® (infliximab)					Maintenance dose: 8-week supply. Refill x 1 year unless noted otherwise.	
☐ Inflectra® (infliximab-dyyb)	☐ 3mg/kg	m	g iv at week: 0, 2, 6		′	
☐ Renflexis® (infliximab-abda)	☐ Other:				Refill x 1 year unless noted otherwise.	
☐ Avsola® (infliximab-axxq	☐ Maintenance dose: (_ mg/kg)	mg IV every	_ weeks	Other:	
Pre-Medication	Dose/Strength			Directi	ons	
☐ Acetaminophen	□ 500mg	☐ Take	1-2 tablets PO prior to inf	usion or p	ost-infusion as directed	
	☐ 25mg IV/PO	☐ Take	1 tablet PO prior to infusion	on or as di	rected OR	
☐ Diphenhydramine			ect contents of 1 vial IV prior to infusion or as directed			
	☐ 40mg 100mg		t contents of 1 vial IV prior			
☐ Methylprednisolone			her: Inject 100mg IV 30 minutes prior to infusion			
INFUSION REACTION ORDE	RS					
Mild reaction protocol:						

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



REMICADE® (infliximab) & Biosimilar

Date:

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

			Copy of The Patient's Up-to-Date Clinical Notes			
☑ Diphenhydramine 25mg IV, one time, for prurit	us.					
If symptoms worsen, see orders for moderate to se	vere reactions.					
Moderate reaction protocol:						
oximes Acetaminophen 650mg PO, one time, for pyrexi	ia or rigors					
☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria						
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms						
If symptoms worsen, see interventions for severe reactions						
Severe reaction protocol: (Call 911 if initiated):						
☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)						
☑ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis						
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis						
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms						
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolate	teral aspect of thigh of ana	phylaxis, may repeat x1	in 5-15 minutes if symptoms are not resolved or			
worsen						
FLUSHING & LOCKING ORDERS						
Flushing Protocol (>66lbs/33kg)						
Flushing Protocol (>66lbs/33kg) PIV and Midline:		Implanted Port, PICC, 1	Tunneled Catheter, and Non-tunneled Catheter:			
, , , , , ,	d after each infusion	•	de 5mL IV flush before infusion/lab draw and 10mL			
PIV and Midline:	d after each infusion	⊠ 0.9% Sodium Chlorid	de 5mL IV flush before infusion/lab draw and 10mL			
PIV and Midline: ☑ 0.9% Sodium Chloride 2-5mL IV flush before and	d after each infusion PICC:	⊠ 0.9% Sodium Chlorid	de 5mL IV flush before infusion/lab draw and 10mL			
PIV and Midline: ☑ 0.9% Sodium Chloride 2-5mL IV flush before and Locking Protocol (>66lbs/33kg) PIV and Midline:		☑ 0.9% Sodium Chlorid IV flush after infusion/l	de 5mL IV flush before infusion/lab draw and 10mL ab draw			
PIV and Midline: ☑ 0.9% Sodium Chloride 2-5mL IV flush before and Locking Protocol (>66lbs/33kg)	PICC:	☑ 0.9% Sodium Chloric IV flush after infusion/I	de 5mL IV flush before infusion/lab draw and 10mL ab draw Implanted Port, Tunneled Catheter, and Non-			
PIV and Midline: ☑ 0.9% Sodium Chloride 2-5mL IV flush before and Locking Protocol (>66lbs/33kg) PIV and Midline: ☑ Heparin Sodium 10 units/mL 1mL IV final	PICC: ⊠ Heparin Sodium 10 ur flush post normal saline	⊠ 0.9% Sodium Chloric IV flush after infusion/l nits/mL 3mL IV final flush	Implanted Port, Tunneled Catheter, and Nontunneled Catheter: Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush			
PIV and Midline:	PICC: ⊠ Heparin Sodium 10 ur flush post normal saline	⊠ 0.9% Sodium Chloric IV flush after infusion/l nits/mL 3mL IV final flush	Implanted Port, Tunneled Catheter, and Nontunneled Catheter: Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush			

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

Prescriber Signature

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.