

## **OCREVUS ZUNOVO™**

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's up-to-Date Clinical Notes

PATIENT INFORMATION (C	Complete or Fax Existing Chart	PRESCRIBER INFORMATION	
Address: City, State, Zip: Phone: Email:	DOB: Alt. Phone: SS#:(lbs) Ht:	State License: Tax ID:	ax:
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)			
Primary Insurance:		Plan #: Group #:	
CLINICAL INFORMATION			
□ G35.A Relapsing-remitting       □ G35.B0 Primary progressive multiple sclerosis       □ G35.B1 Active primary progressive multiple sclerosis       □ G35.B2 Non-active primary progressive multiple sclerosis         □ G35.C0 Secondary progressive multiple sclerosis, unspecified       □ G35.C1 Active secondary progressive multiple sclerosis       □ G35.D Multiple sclerosis, multiple sclerosis         □ Other (Specify ICD-10 Code):       □ Past DMT Therapies:         □ Date of Last MRI:       Past DMT Therapies:         □ Date of last treatment with an OCREVUS product (if applicable) (MM/DD/YYYY):       □ Quantitative Serum Immunoglobulins Test Results:			
ORDERS			
Prescription type: ☐ New start  Medication	☐ Restart ☐ Continued therapy Total Doses Received: Date of Last Injection/Infusion:  Dose/Frequency Quantity/Refills		
☐ OCREVUS ZUNOVO™ (920mg/23mL)	☐ 920 mg/23,000 units (920 mg ocrelizumab and 23,000 units of hyaluronidase) administered as a single 23 mL subcutaneous injection in the abdomen over approximately 10 minutes every 6 months ☐ Other:		☐ Quantity: 1 Vial
Pre-Medication	Dose/Strength	Directions	
☐ Acetaminophen	□ 500mg	☐ Take 1-2 tablets PO prior to infusion or post-infusion as directed	
☐ Cetirizine	□ 10mg	☐ Take 1 tablet PO prior to infusion or as directed	
☐ Dexamethasone	☐ 20mg	☐ Take 1 tablet PO prior to infusion or as directed	
INFUSION REACTION ORDERS			
Mild reaction protocol:  ☑ Diphenhydramine 25mg IV, or  If symptoms worsen, see orders for the symptoms worsen.	· ·		

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



☐ Acetaminophen 650mg PO, one time, for pyrexia or rigors

## **OCREVUS ZUNOVO™**

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's up-to-Date Clinical Notes

<ul> <li>☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms</li> <li>If symptoms worsen, see interventions for severe reactions</li> <li>Severe reaction protocol: (Call 911 if initiated):</li> <li>☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (&gt;95%)</li> </ul>				
Severe reaction protocol: (Call 911 if initiated):				
☐ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)				
☑ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis				
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis				
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms				
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or				
worsen				
FLUSHING & LOCKING ORDERS				
Flushing Protocol (>66lbs/33kg)				
PIV and Midline: Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:				
□ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion □ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10m				
IV flush after infusion/lab draw				
Locking Protocol (>66lbs/33kg)				
PIV and Midline: PICC: Implanted Port, Tunneled Catheter, and Non-				
☐ Heparin Sodium 10 units/mL 1mL IV final ☐ Heparin Sodium 10 units/mL 3mL IV final ☐ tunneled Catheter:				
flush post normal saline flush  flush post normal saline flush  ☐ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush				
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused				
SIGNATURE				
We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.				
X Date:				
Prescriber Signature				

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.