

## **ULTOMIRIS®**

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

PATIENT INFORMATION (Complete or Fax Existing Chart)				PRESCRIBER INFORMATION				
Name:				Prescriber Name:				
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)								
Primary Insurance:			Plan #:					
CLINICAL INFORMATION								
Primary ICD-10 Code (Please Specify Diagnosis):								
ULTOMIRIS® ORDERS								
Prescription type:   New start   Restart   Continued therapy Total Doses Received:   Date of Last Injection/Infusion:						ion:		
Medication  Intravenous Ultomiris® (ravulizumab)	Strength  ☐ 1,100mg/11mL vial ☐ 300mg/3mL vial ☐ 300mg/30mL vial ☐ Other:		Dose/Frequency  Loading dose: Begin mg IV on day 1  Then 2 weeks later  Maintenance dose: Begin mg IV every weeks  Other:					
Subcutaneous Ultomiris® (ravulizumab)	245mg/3.5 mL prefilled cartridge with on body injector		□ 490 mg once weekly in adult patients greater than or equal to 40 kg body weight with PNH or aHUS.  □ Other:					
Pre-Medication		Dose/Strength		Directions				
☐ Acetaminophen		□ 500mg		☐ Take 1-2 tablets PO prior to infusion or post-infusion as directed				
☐ Diphenhydramine		☐ 25mg IV/PO ☐ 50mg IV/PO		☐ Take 1 tablet PO prior to infusion or as directed OR ☐ Inject contents of 1 vial IV prior to infusion or as directed				
☐ Methylprednisolone		□ 40mg □ 125mg		☐ Inject contents of 1 vial IV prior to infusion or as directed				
INFLISION REACTION ORDERS								

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



## **ULTOMIRIS®**

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

Date:

		Please Sen	Please Send a Copy of The Patient's Up-to-Date Clinical Note		
Mild reaction protocol:					
☑ Diphenhydramine 25mg IV, one time, for pru	ritus.				
If symptoms worsen, see orders for moderate to	severe reactions.				
Moderate reaction protocol:					
oximes Acetaminophen 650mg PO, one time, for pyro	exia or rigors				
☑ Diphenhydramine 50mg IV, one time, for pru	ritus or urticaria				
oximes Methylprednisolone 125mg IV, one time, for	respiratory or neurologic sy	mptoms			
If symptoms worsen, see interventions for severe	e reactions				
Severe reaction protocol: (Call 911 if initiated):					
$oximes$ Titrate oxygen via continuous flow per nasal ${f c}$	cannula or face mask to ma	intain spO2 of greater th	an ninety-five percent (>95%)		
☑ Diphenhydramine 50mg IV,one time, for resp	iratory symptoms, edema,	or anaphylaxis			
☑ Methylprednisolone 125mg IV, one time, for	respiratory symptoms, eder	ma, or anaphylaxis			
⊠ Sodium Chloride 0.9% 500mL IV over 30-60 m	nin, one time, for cardiovasc	cular symptoms			
☑ Epinephrine 0.3mg/0.3mL IM into mis-antero	lateral aspect of thigh of an	aphylaxis, may repeat x1	l in 5-15 minutes if symptoms are not resolved or		
worsen					
FLUSHING & LOCKING ORDERS					
Flushing Protocol (>66lbs/33kg)					
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:			
⊠ 0.9% Sodium Chloride 2-5mL IV flush before a	and after each infusion	☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL			
		IV flush after infusion/lab draw			
Locking Protocol (>66lbs/33kg)					
PIV and Midline:	d Midline: PICC:		Implanted Port, Tunneled Catheter, and Non-		
☐ Heparin Sodium 10 units/mL 1mL IV final	eparin Sodium 10 units/mL 1mL IV final		tunneled Catheter:		
flush post normal saline flush	flush post normal saline	e flush	⋈ Heparin Sodium 100 units/mL 3-5mL IV final		
			flush post normal saline flush		
** May substitute Dextrose 5% in Water, or alternati	ve, for 0.9& Sodium Chloride,	when indicated due to inco	mpatibility with medications bring infused		
SIGNATURE					
We hereby authorize Talis Healthcare LLC to pro medicine as prescribed in this referral.	vide all supplies and additic	onal services (nursing/pa	tient training) required to provide and deliver the		

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

**Prescriber Signature** 

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.